


Final Exam — Dupuy — Math 121 — Fall 2016

**Instructions** Remember to show all your work to get full credit. Please leave answers in their exact form. This is a closed book test. You may not use a calculator. If you need extra paper let me know. You will be marked off for floating expressions and equalities not related to the problem. You can potentially be marked off for being vague or imprecise.

Name: **KEY**  
Section: 121B

Problem	Possible	Score
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
11	10	
12	10	
EC1	10	
EC2	10	
Total	120	
Percentage		

1. Make a level set plot of the function  $f(x, y) = x^2 + y^2$ .



2. Compute the following directions derivatives. If the direction is not a unit vector, normalize it.

(a)  $(D_{\vec{u}}f)(1,1)$  where  $f(x,y) = x^2 + y^2$  and  $\vec{u} = (1,0)$ .

(b)  $(D_{\vec{v}}g)(0,0,0)$  where  $g(x,y,z) = ze^{xy}$  and  $\vec{v} = (1,1,1)$ .

(a)  $\nabla f = (2x, 2y), \nabla f(1,1) = (2,2)$   
 $(D_{\vec{u}}f)(1,1) = \nabla f(1,1) \cdot (1,0) = (2,2) \cdot (1,0) = 2.$

(b)  ~~$(D_{\vec{v}}g)(0,0,0)$~~   $\nabla g = (yze^{xy}, xze^{xy}, e^{xy})$   
 $\nabla g(0,0,0) = (0,0,1).$

~~$(D_{\vec{v}}g)(0,0,0)$~~   $D_{\vec{v}}g(0,0,0) = \nabla g(0,0,0) \cdot \frac{\vec{v}}{|\vec{v}|}$   
 $= (0,0,1) \cdot \frac{(1,1,1)}{\sqrt{3}}$   
 $= \frac{1}{\sqrt{3}}.$

3. Find a parametrization of the line passing through the points  $(1, 2, 0)$  and  $(0, 1, 1)$ .

$$\begin{aligned}\vec{r}(t) &= (1, 2, 0)t + (0, 1, 1)(1-t) \\ &= (t, 2t, 0) + (0, 1-t, 1-t) \\ &= (t, 2t + (1-t), 0 + (1-t)) \\ &= (t, 1+t, 1-t).\end{aligned}$$

4. Let  $\vec{a} = (1, 1, 0)$  and  $\vec{b} = (-1, 1, 1)$

(a) Compute  $\vec{a} \cdot \vec{b}$ .

(b) Find the angle between  $\vec{a}$  and  $\vec{b}$ .

$$(a) \quad \vec{a} \cdot \vec{b} = (1, 1, 0) \cdot (-1, 1, 1) \\ = 0.$$

$$(b) \quad 90^\circ.$$

5. Find the line tangent to the curve  $\alpha(t) = (e^t, e^{2t}, e^{3t})$  at the point  $(1, 1, 1)$ .

$$\vec{\alpha}(0) = (1, 1, 1)$$

$$\vec{\alpha}'(t) = (e^t, 2e^{2t}, 3e^{3t})$$

$$\vec{\alpha}'(0) = (1, 2, 3)$$

$$\vec{q}(t) = (1, 1, 1) + t(1, 2, 3).$$

6. Find the plane tangent to the surface  $x^2 + y^3 + z^4 = 3$  at the point  $(1, 1, 1)$ .

$$\nabla g = (2x, 3y, 4z)$$

$$\nabla g(1, 1, 1) = (2, 3, 4) = \vec{n} \quad \vec{r}_0 = (1, 1, 1)$$

Plane eqn:  $\vec{n} \cdot (\vec{r} - \vec{r}_0) = 0.$

$$\Rightarrow (2, 3, 4) \cdot (x-1, y-1, z-1) = 0$$

$$\Rightarrow 2(x-1) + 3(y-1) + 4(z-1) = 0.$$

7. Find an equation of the plane containing the lines  $\vec{l}_1(t) = (2t + 1, t, 0)$  and  $\vec{l}_2(t) = (1, 3t, 2t)$ . (These lines intersect btw)

$$\vec{v}_1 = (2, 1, 0)$$

$$\vec{v}_2 = (0, 3, 2)$$

$$\vec{n} = \vec{v}_1 \times \vec{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & 0 \\ 0 & 3 & 2 \end{vmatrix}$$

$$= \mathbf{i}(2) - \mathbf{j}(4) + \mathbf{k}(6)$$

$$= (2, -4, 6)$$

pt of intersection:

$$\vec{l}_1(t) = \vec{l}_2(\tau)$$

$$\Leftrightarrow \begin{cases} 2t+1 = 1 \\ t = 3\tau \\ 0 = 2\tau \end{cases}$$

$$\Rightarrow \tau = 0 \Rightarrow t = 0$$

pt of intersection:  
(1, 0, 0)

$$2(x-1) - 4y + 6z = 0$$

$$\Rightarrow 2x - 2 - 4y + 6z = 0$$

$$\Rightarrow 2x - 4y + 6z = 2$$

$$\Leftrightarrow \boxed{x - 2y + 3z = 1}$$



8. Find and classify the critical points of the function  $f(x, y) = x^2 - 2x + 1 - y^2$ .

$$\begin{cases} f_x = 2x - 2 = 0 \\ f_y = -2y = 0 \end{cases}$$

$$\Rightarrow \begin{matrix} x=1, y=0 \\ \text{only CP is} \\ (1, 0). \end{matrix}$$


2nd Der test:

$$\begin{aligned} H(1, 0) &= \det \begin{bmatrix} f_{xx}(1, 0) & f_{xy}(1, 0) \\ f_{yx}(1, 0) & f_{yy}(1, 0) \end{bmatrix} \\ &= \det \begin{bmatrix} 2 & 0 \\ 0 & -2 \end{bmatrix} = -4 < 0 \end{aligned}$$

$\Rightarrow$  saddle point. //

9. Compute the volume of the region  $E = \{(x, y, z) : 0 \leq z \leq 1 - x^2 - y^2\}$ .

$$\begin{cases} z \in [0, 1 - x^2 - y^2] = [0, 1 - r^2] \\ r \in [0, 1] \\ \theta \in [0, 2\pi] \end{cases}$$



$$\iiint_E dV = \int_0^{2\pi} \int_0^1 \int_0^{1-r^2} r \, dz \, dr \, d\theta$$

$$= \int_0^{2\pi} \int_0^1 r(1-r^2) \, dr \, d\theta$$

$$= \left( \int_0^{2\pi} d\theta \right) \left( \int_0^1 (r - r^3) \, dr \right)$$



$$= 2\pi \left( \frac{1}{2} - \frac{1}{4} \right) = 2\pi \left( \frac{1}{4} \right) = \frac{\pi}{2} //$$

10. Let  $E$  be the region below the  $xy$ -plane and above the paraboloid  $z + 1 = x^2 + y^2$ .  
 Compute

$$\iiint_E z dV.$$

$$\begin{aligned} & \iiint z dV \\ &= \int_0^{2\pi} \int_0^1 \int_{r^2-1}^0 z r dz dr d\theta \\ &= \int_0^{2\pi} \int_0^1 \frac{(r^2-1)^3}{2} r dr d\theta \\ &= \left( \int_0^{2\pi} d\theta \right) \left( \int_0^1 \frac{(r^2-1)^2}{2} r dr \right) \\ &= -2\pi \left( \int_0^1 \frac{u^2}{4} du \right) \\ &= -\frac{2\pi}{4} \left( \int_{-1}^0 u^2 du \right) \\ &= -\frac{\pi}{2} \left( \frac{u^3}{3} \Big|_{u=-1}^{u=0} \right) \end{aligned}$$

$$= -\left(\frac{\pi}{2}\right) \left( \frac{0}{3} - \left( \frac{(-1)^3}{3} \right) \right) = -\frac{\pi}{2} \left( \frac{1}{3} \right) = -\frac{\pi}{6}$$




$$\begin{aligned} u &= r^2 - 1 \\ du &= 2r dr \\ \Rightarrow \frac{du}{2} &= r dr \\ r=0 &\Rightarrow u=-1 \\ r=1 &\Rightarrow u=0 \end{aligned}$$

11. Let  $M = \{(x, y, z) : x^2 + y^2 + z^2 e^z = e^z \text{ \& } z \geq 0\}$  with an upward pointing normal (this surface looks essentially like the upper half sphere). Compute

$$\iint_M \nabla \times \mathbf{F} \cdot d\mathbf{S}$$

Where  $\mathbf{F} = (-y, x, z)$ .

Stokes:  $\iint_M \nabla \times \mathbf{F} \cdot d\mathbf{S} = \int_{\partial M} \mathbf{F} \cdot d\mathbf{r}$



(on the plane)  
 $z=0$

$$= \int_{\partial M} -y dx + x dy + z dz \rightarrow 0$$

$$= \int_{\partial M} -y dx + x dy$$

$$= \iint_D \left[ \frac{\partial}{\partial x}(x) - \frac{\partial}{\partial y}(-y) \right] dA$$

$$= 2 \iint_D dA = 2\pi(1)^2 = 2\pi //$$

12. Compute the flux of  $\mathbf{F} = (x, y, z)$  through the boundary of the tetrahedron  $T$  given by

$$T = \{(x, y, z) : x + y + z \leq 2 \text{ \& } x \geq 0 \text{ \& } y \geq 0 \text{ \& } z \geq 0\}$$

Divergence theorem

$$\iint_{\partial T} \mathbf{F} \cdot \mathbf{N} \, dS = \iiint_T (\nabla \cdot \mathbf{F}) \, dV$$

$$= \iiint_T 3 \, dV$$


$$= 3 \int_0^2 \int_0^{2-x} \int_0^{2-x-y} dz \, dy \, dx$$

$$= 3 \int_0^2 \int_0^{2-x} (2-x-y) \, dy \, dx$$

$$= 3 \int_0^2 \left[ (2-x)(2-x) - \frac{(2-x)^2}{2} \right] dx$$

$$= 3 \int_0^2 \frac{(2-x)^2}{2} dx$$

$$= \frac{3}{2} \int_0^2 \frac{d}{dx} \left[ -\frac{(2-x)^3}{3} \right] dx = \frac{3}{2} \left[ \frac{8}{3} \right] = 4.$$



**EC1** Let  $M$  be a simple closed oriented surface in  $\mathbf{R}^3$ . Prove Stoke's Theorem:

$$\iint_M \nabla \times \mathbf{F} \cdot d\mathbf{S} = \oint_{\partial M} \mathbf{F} \cdot d\mathbf{r}.$$

**EC2** Compute  $\int_{-\infty}^{\infty} e^{-x^2} dx$ . (Hint: this computation involved a double integral and a change to polar coordinates.)